

## ARG65669 anti-Annexin A2 antibody

Package: 100 µg  
Store at: -20°C

### Summary

Product Description	Goat Polyclonal antibody recognizes Annexin A2
Tested Reactivity	Hu, Ms
Predict Reactivity	Cow, Rat, Pig
Tested Application	WB
Specificity	This antibody is expected to recognise both reported isoforms. NP_004030.1, NP_001002857.1 and NP_001129487.1 are variants of isoform 2.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	Annexin A2
Species	Human
Immunogen	Synthetic peptide around the N-terminus of Human Annexin A2 (STVHEILCKLSLEGD)
Conjugation	Un-conjugated
Alternate Names	Placental anticoagulant protein IV; LPC2; Protein I; Calpactin-1 heavy chain; Calpactin I heavy chain; p36; CAL1H; ANX2; LPC2D; Lipocortin II; ANX2L4; HEL-S-270; LIP2; Annexin II; Annexin A2; Annexin-2; P36; Chromobindin-8; PAP-IV

### Application Instructions

Application table	Application	Dilution
	WB	0.5 - 3 µg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated

freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Database links	<a href="#">GeneID: 12306 Mouse</a> <a href="#">GeneID: 302 Human</a> <a href="#">Swiss-port # P07355 Human</a> <a href="#">Swiss-port # P07356 Mouse</a>
Gene Symbol	ANXA2
Gene Full Name	annexin A2
Background	This gene encodes a member of the annexin family. Members of this calcium-dependent phospholipid-binding protein family play a role in the regulation of cellular growth and in signal transduction pathways. This protein functions as an autocrine factor which heightens osteoclast formation and bone resorption. This gene has three pseudogenes located on chromosomes 4, 9 and 10, respectively. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]
Function	Calcium-regulated membrane-binding protein whose affinity for calcium is greatly enhanced by anionic phospholipids. It binds two calcium ions with high affinity. May be involved in heat-stress response. Inhibits PCSK9-enhanced LDLR degradation, probably reduces PCSK9 protein levels via a translational mechanism but also competes with LDLR for binding with PCSK9 [UniProt]
Research Area	Signaling Transduction antibody
Calculated Mw	39 kDa
PTM	Phosphorylation of Tyr-24 enhances heat stress-induced translocation to the cell surface. ISGylated.

## Images



ARG65669 anti-Annexin A2 antibody WB image

Western Blot: A431 lysate (RIPA buffer, 35 µg total protein per lane) stained with ARG65669 anti-Annexin A2 antibody at 0.5 µg/ml dilution.