

## ARG66864 anti-NSE / Neuron Specific Enolase antibody

Package: 100 μg Store at: -20°C

## Summary

duct DescriptionRabbit Polyclonal antibody recognizes NSE / Neuron Specific Enolaseted ReactivityHudict ReactivityMs, Ratted ApplicationICC/IF, IHC-P, WB
dict Reactivity Ms, Rat
ted Application ICC/IF, IHC-P, WB
t Rabbit
nality Polyclonal
ype IgG
get Name NSE / Neuron Specific Enolase
cies Human
Synthetic peptide corresponding to aa. 371-420 of Human NSE / Neuron Specific Enolase.
jugation Un-conjugated
rnate Names Neural enolase; NSE; Enolase 2; Gamma-enolase; 2-phospho-D-glycerate hydro-lyase; HEL-S-279; Neuron-specific enolase; EC 4.2.1.11

## **Application Instructions**

Application table	Application	Dilution
	ICC/IF	1:200 - 1:1000
	IHC-P	1:100 - 1:300
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HepG2	
Observed Size	~ 48 kDa	

### Properties

Form	Liquid	
Purification	Affinity purification with immunogen.	
Buffer	PBS, 0.02% Sodium azide, 50% Glycerol and 0.5% BSA.	
Preservative	0.02% Sodium azide	
Stabilizer	50% Glycerol and 0.5% BSA	
Concentration	1 mg/ml	

Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Gene Symbol	ENO2
Gene Full Name	enolase 2 (gamma, neuronal)
Background	This gene encodes one of the three enolase isoenzymes found in mammals. This isoenzyme, a homodimer, is found in mature neurons and cells of neuronal origin. A switch from alpha enolase to gamma enolase occurs in neural tissue during development in rats and primates. [provided by RefSeq, Jul 2008]
Function	Has neurotrophic and neuroprotective properties on a broad spectrum of central nervous system (CNS) neurons. Binds, in a calcium-dependent manner, to cultured neocortical neurons and promotes cell survival (By similarity). [UniProt]
Calculated Mw	47 kDa
Cellular Localization	Cytoplasm. Cell membrane. Note=Can translocate to the plasma membrane in either the homodimeric (alpha/alpha) or heterodimeric (alpha/gamma) form. [UniProt]

#### Images



#### ARG66864 anti-NSE / Neuron Specific Enolase antibody ICC/IF image

Immunofluorescence: HeLa cells stained with ARG66864 anti-NSE / Neuron Specific Enolase antibody (green) at 1:200 dilution, overnight at 4°C. DAPI (blue) for nuclear staining.



#### ARG66864 anti-NSE / Neuron Specific Enolase antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human brain tissue stained with ARG66864 anti-NSE / Neuron Specific Enolase antibody. The picture on the right is blocked with the synthetic peptide.

# - 117 - 85 - 48 - 34 - 26 - 19 HepG2



Western blot: HepG2 cell lysate stained with ARG66864 anti-NSE / Neuron Specific Enolase antibody at 1:2000 dilution.



#### ARG66864 anti-NSE / Neuron Specific Enolase antibody ICC/IF image

Immunofluorescence: A549 cells stained with ARG66864 anti-NSE / Neuron Specific Enolase antibody. The picture on the right is blocked with the synthetic peptide.