

## ARG67274 anti-PIK3CA / p110 alpha (H1047R mutation) antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody recognizes PIK3CA / p110 alpha (H1047R mutation) antibody
Tested Reactivity	Hu
Tested Application	IHC-P, WB
Specificity	The antibody can specifically recognize the h1047r mutation of human PIK3CA protein, and does not react with wild-type PIK3CA protein.
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b, kappa
Target Name	PIK3CA / p110 alpha
Species	Human
Immunogen	Synthetic peptide derived from Human PIK3CA / p110 alpha (H1047R mutation, aa sequence: 1000-1068).
Conjugation	Un-conjugated
Alternate Names	MCM; MCMTC; p110alpha; PI3-kinase subunit alpha; PI3K; CWS5; p110-alpha; PI3Kalpha; CLOVE; EC 2.7.1.153; MCAP; PtdIns-3-kinase subunit p110-alpha; EC 2.7.11.1; Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform; Phosphatidylinositol 4,5-bisphosphate 3-kinase 110 kDa catalytic subunit alpha; PI3K-alpha; PtdIns-3-kinase subunit alpha; Phosphoinositide-3-kinase catalytic alpha polypeptide; Serine/threonine protein kinase PIK3CA

### Application Instructions

Application table	Application	Dilution
	IHC-P	1:200 - 1:1000
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Protein G purification
Buffer	PBS, 0.05% Proclin 300, 50% Glycerol and 0.05% BSA.
Preservative	0.05% Proclin 300
Stabilizer	50% Glycerol and 0.05% BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw

cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


#### Note

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	PIK3CA
Gene Full Name	phosphatidylinositol-4,5-bisphosphate 3-kinase, catalytic subunit alpha
Background	Phosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. The protein encoded by this gene represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has been found to be oncogenic and has been implicated in cervical cancers. [provided by RefSeq, Jul 2008]
Function	Phosphoinositide-3-kinase (PI3K) that phosphorylates PtdIns (Phosphatidylinositol), PtdIns4P (Phosphatidylinositol 4-phosphate) and PtdIns(4,5)P2 (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3). PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Participates in cellular signaling in response to various growth factors. Involved in the activation of AKT1 upon stimulation by receptor tyrosine kinases ligands such as EGF, insulin, IGF1, VEGFA and PDGF. Involved in signaling via insulin-receptor substrate (IRS) proteins. Essential in endothelial cell migration during vascular development through VEGFA signaling, possibly by regulating RhoA activity. Required for lymphatic vasculature development, possibly by binding to RAS and by activation by EGF and FGF2, but not by PDGF. Regulates invadopodia formation in breast cancer cells through the PDK1-AKT1 pathway. Participates in cardiomyogenesis in embryonic stem cells through a AKT1 pathway. Participates in vasculogenesis in embryonic stem cells through PDK1 and protein kinase C pathway. Has also serine-protein kinase activity: phosphorylates PIK3R1 (p85alpha regulatory subunit), EIF4EBP1 and HRAS. [UniProt]
Calculated Mw	124 kDa

## Images



ARG67274 anti-PIK3CA / p110 alpha (H1047R mutation) antibody  
IHC-P image

Immunohistochemistry: Human gastric adenocarcinoma stained with ARG67274 anti-PIK3CA / p110 alpha (H1047R mutation) antibody at dilution.



ARG67274 anti-PIK3CA / p110 alpha (H1047R mutation) antibody  
WB image

Western blot: K562 stained with ARG67274 anti-PIK3CA / p110 alpha (H1047R mutation) antibody at dilution.