

ARG70204 Pig IL4 recombinant protein (Active) (His-tagged, C-ter)

Package: 100 µg, 20 µg
Store at: -20°C

Summary

Product Description	E. coli expressed, His-tagged (C-ter) Active Pig IL4 recombinant protein
Tested Application	SDS-PAGE
Target Name	IL4
Species	Pig
A.A. Sequence	His25 - Cys133
Expression System	E. coli
Activity	Active
Activity Note	Determined by its ability to induce proliferation in TF-1 cells. The ED50 for this effect is < 0.6 ng/mL.
Alternate Names	BSF-1; IL-4; B-cell stimulatory factor 1; BCGF1; Lymphocyte stimulatory factor 1; BCGF-1; Interleukin-4; Pitracinra; Binetrakin; BSF1

Properties


Form	Powder
Purification Note	Endotoxin level is less than 0.1 EU/µg of the protein, as determined by the LAL test.
Purity	> 98% (by SDS-PAGE)
Buffer	PBS (pH 7.4)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.
Storage instruction	For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	IL4
Gene Full Name	interleukin 4
Background	The protein encoded by this gene is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

Function	Participates in at least several B-cell activation processes as well as of other cell types. It is a costimulator of DNA-synthesis. It induces the expression of class II MHC molecules on resting B-cells. It enhances both secretion and cell surface expression of IgE and IgG1. It also regulates the expression of the low affinity Fc receptor for IgE (CD23) on both lymphocytes and monocytes. [UniProt]
Cellular Localization	Secreted. [UniProt]

Images



ARG70204 Pig IL4 recombinant protein (Active) (His-tagged, C-ter)
SDS-PAGE image

SDS-PAGE analysis of ARG70204 Pig IL4 recombinant protein (Active) (His-tagged, C-ter).