

Product datasheet

info@arigobio.com

ARG70212
Pig GM-CSF recombinant protein (Active) (His-tagged, N-ter)

Package: 100 μg, 20 μg

Store at: -20°C

Summary

Product Description E. coli expressed, His-tagged (N-ter) Active Pig GM-CSF recombinant protein

Tested Application SDS-PAGE
Target Name GM-CSF

Species Pig

A.A. Sequence Ala18 - Lys144

Expression System E. coli
Activity Active

Activity Note Determined by its ability to induce proliferation in TF-1 cells. The ED50 for this effect is < 3 ng/mL.

Alternate Names GM-CSF; Granulocyte-macrophage colony-stimulating factor; CSF; GMCSF; Molgramostin; Colony-

stimulating factor; Sargramostim

Properties

Form Powder

Purification Note Endotoxin level is less than 0.1 EU/μg of the protein, as determined by the LAL test.

Purity > 98% (by SDS-PAGE)

Buffer PBS (pH 7.4)

Reconstitution It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less

than 200 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure

the protein is dissolved completely.

Storage instruction For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and

store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid

repeated freeze/thaw cycles. Suggest spin the vial prior to opening.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol CSF2

Gene Full Name colony stimulating factor 2 (granulocyte-macrophage)

Background The protein encoded by this gene is a cytokine that controls the production, differentiation, and

function of granulocytes and macrophages. The active form of the protein is found extracellularly as a homodimer. This gene has been localized to a cluster of related genes at chromosome region 5q31, which is known to be associated with interstitial deletions in the 5q- syndrome and acute myelogenous leukemia. Other genes in the cluster include those encoding interleukins 4, 5, and 13. [provided by

RefSeq, Jul 2008]

Function Cytokine that stimulates the growth and differentiation of hematopoietic precursor cells from various

lineages, including granulocytes, macrophages, eosinophils and erythrocytes. [UniProt]

Highlight Related products:

GM-CSF antibodies; GM-CSF ELISA Kits; GM-CSF Duos / Panels; GM-CSF recombinant proteins;


Related news:

HMGB1 in inflammation Inflammatory Cytokines

Cellular Localization

Secreted. [UniProt]

Images



ARG70212 Pig GM-CSF recombinant protein (Active) (His-tagged, Nter) SDS-PAGE image

SDS-PAGE analysis of ARG70212 Pig GM-CSF recombinant protein (Active) (His-tagged, N-ter).

2/2