

ARG70276 Human MMP1 recombinant protein (Active) (His-tagged, C-ter)

Package: 50 μg Store at: -20°C

Summary

Fested ReactivityHuFested ApplicationFuncSt, SDS-PAGEFarget NameMMP1SpeciesHumanA.A. SequencePhe20 - Asn469 of Human MMP1 (NP_002412.1) with 6X His tag at the C - terminus.Expression SystemHEK293ActivityActiveActivity NoteMeasured in a cell migration assay using A549 cells. 1 ng/ml of Recombinant Human MMP-1 can effectively induce A549 cells migration.		
Fested ApplicationFuncSt, SDS-PAGEFarget NameMMP1SpeciesHumanA.A. SequencePhe20 - Asn469 of Human MMP1 (NP_002412.1) with 6X His tag at the C - terminus.Expression SystemHEK293ActivityActiveActivity NoteMeasured in a cell migration assay using A549 cells. 1 ng/ml of Recombinant Human MMP-1 can effectively induce A549 cells migration.Alternate NamesMMP-1; CLG; Fibroblast collagenase; Matrix metalloproteinase-1; CLGN; EC 3.4.24.7; Interstitial	Product Description	HEK293 expressed, His-tagged (C-ter) Active Human MMP1 recombinant protein.
Target NameMMP1SpeciesHumanA.A. SequencePhe20 - Asn469 of Human MMP1 (NP_002412.1) with 6X His tag at the C - terminus.Expression SystemHEK293ActivityActiveActivity NoteMeasured in a cell migration assay using A549 cells. 1 ng/ml of Recombinant Human MMP-1 can effectively induce A549 cells migration.Alternate NamesMMP-1; CLG; Fibroblast collagenase; Matrix metalloproteinase-1; CLGN; EC 3.4.24.7; Interstitial	Tested Reactivity	Hu
SpeciesHumanA.A. SequencePhe20 - Asn469 of Human MMP1 (NP_002412.1) with 6X His tag at the C - terminus.Expression SystemHEK293ActivityActiveActivity NoteMeasured in a cell migration assay using A549 cells. 1 ng/ml of Recombinant Human MMP-1 can effectively induce A549 cells migration.Alternate NamesMMP-1; CLG; Fibroblast collagenase; Matrix metalloproteinase-1; CLGN; EC 3.4.24.7; Interstitial	Tested Application	FuncSt, SDS-PAGE
A.A. SequencePhe20 - Asn469 of Human MMP1 (NP_002412.1) with 6X His tag at the C - terminus.Expression SystemHEK293ActivityActiveActivity NoteMeasured in a cell migration assay using A549 cells. 1 ng/ml of Recombinant Human MMP-1 can effectively induce A549 cells migration.Alternate NamesMMP-1; CLG; Fibroblast collagenase; Matrix metalloproteinase-1; CLGN; EC 3.4.24.7; Interstitial	Target Name	MMP1
Expression SystemHEK293ActivityActiveActivity NoteMeasured in a cell migration assay using A549 cells. 1 ng/ml of Recombinant Human MMP-1 can effectively induce A549 cells migration.Alternate NamesMMP-1; CLG; Fibroblast collagenase; Matrix metalloproteinase-1; CLGN; EC 3.4.24.7; Interstitial	Species	Human
ActivityActiveActivity NoteMeasured in a cell migration assay using A549 cells. 1 ng/ml of Recombinant Human MMP-1 can effectively induce A549 cells migration.Alternate NamesMMP-1; CLG; Fibroblast collagenase; Matrix metalloproteinase-1; CLGN; EC 3.4.24.7; Interstitial	A.A. Sequence	Phe20 - Asn469 of Human MMP1 (NP_002412.1) with 6X His tag at the C - terminus.
Activity NoteMeasured in a cell migration assay using A549 cells. 1 ng/ml of Recombinant Human MMP-1 can effectively induce A549 cells migration.Alternate NamesMMP-1; CLG; Fibroblast collagenase; Matrix metalloproteinase-1; CLGN; EC 3.4.24.7; Interstitial	Expression System	HEK293
effectively induce A549 cells migration.Alternate NamesMMP-1; CLG; Fibroblast collagenase; Matrix metalloproteinase-1; CLGN; EC 3.4.24.7; Interstitial	Activity	Active
	Activity Note	
	Alternate Names	

Properties

Form	Powder
Purification Note	0.22 μm filter sterilized. Endotoxin level is 97% (by SDS-PAGE)
Buffer	PBS (pH 7.4)
Reconstitution	Reconstitute to a concentration of 0.1 - 0.5 mg/ml in sterile distilled water.
Storage instruction	For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and store at -20°C for up to one month, at 2-8°C for up to one week. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	MMP1
Gene Full Name	matrix metallopeptidase 1
Function	Cleaves collagens of types I, II, and III at one site in the helical domain. Also cleaves collagens of types VII and X (PubMed:2557822, PubMed:2153297, PubMed:1645757). In case of HIV infection, interacts and cleaves the secreted viral Tat protein, leading to a decrease in neuronal Tat's mediated neurotoxicity (PubMed:16807369). [UniProt]
Calculated Mw	54 kDa
РТМ	Undergoes autolytic cleavage to two major forms (22 kDa and 27 kDa). A minor form (25 kDa) is the glycosylated form of the 22 kDa form. The 27 kDa form has no activity while the 22/25 kDa form can act as activator for collagenase.

Tyrosine phosphorylated in platelets by PKDCC/VLK. [UniProt] **Cellular Localization** Secreted, extracellular space, extracellular matrix. [UniProt] Images ARG70276 Human MMP1 recombinant protein (Active) (His-tagged, - 135 C-ter) SDS-PAGE image - 98 - 72 SDS-PAGE analysis of ARG70276 Human MMP1 recombinant protein - 55 (Active) (His-tagged, C-ter). - 42 - 36 - 24 - 16 Human MMP1 recombinant protein (His-tagged, C-ter)