

ARG70469 Human CD70 recombinant protein (Active) (His-tagged, N-ter)

Package: 100 µg, 20 µg
Store at: -20°C

Summary

Product Description	E. coli expressed, His-tagged (N-ter) Active Human CD70 recombinant protein
Tested Application	SDS-PAGE
Target Name	CD70
Species	Human
A.A. Sequence	Gln39 - Pro193
Expression System	E. coli
Activity	Active
Activity Note	Determined by its ability to induce IL-8 secretion in human PBMCs. The ED50 for this effect is < 0.6 ng/mL.
Alternate Names	CD70; CD70 Molecule; CD27L; CD70 Antigen; CD27 Ligand; CD27-L; CD27LG; TNFSF7; Tumor Necrosis Factor Ligand Superfamily Member 7; Tumor Necrosis Factor (Ligand) Superfamily, Member 7

Properties

Form	Powder
Purification Note	Endotoxin level is less than 0.1 EU/µg of the protein, as determined by the LAL test.
Purity	> 98% (by SDS-PAGE)
Buffer	PBS containing 0.05% sarkosyl (pH 8.0)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.
Storage instruction	For long term, lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	CD70
Gene Full Name	CD70 Molecule
Background	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis.
Function	Cytokine which is the ligand for CD27. The CD70-CD27 pathway plays an important role in the generation and maintenance of T cell immunity, in particular during antiviral responses. Upon CD27 binding, induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-

cells.


PTM

Disulfide bond, Glycoprotein

Cellular Localization

Membrane

Images



ARG70469 Human CD70 recombinant protein (Active) (His-tagged, N-ter) SDS-PAGE image

SDS-PAGE analysis of ARG70469 Human CD70 recombinant protein (Active) (His-tagged, N-ter)