

ARG81557 Human ADAMTS1 ELISA Kit

Package: 96 wells
Store at: 4°C

Component

Cat. No.	Component Name	Package	Temp
ARG81557-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG81557-002	Standard	2 X 10 ng/vial	4°C
ARG81557-003	Standard/Sample diluent	30 ml (Ready to use)	4°C
ARG81557-004	Antibody conjugate concentrate (100X)	1 vial (100 µl)	4°C
ARG81557-005	Antibody diluent buffer	12 ml (Ready to use)	4°C
ARG81557-006	HRP-Streptavidin concentrate (100X)	1 vial (100 µl)	4°C
ARG81557-007	HRP-Streptavidin diluent buffer	12 ml (Ready to use)	4°C
ARG81557-008	25X Wash buffer	20 ml	4°C
ARG81557-009	TMB substrate	10 ml (Ready to use)	4°C (Protect from light)
ARG81557-010	STOP solution	10 ml (Ready to use)	4°C
ARG81557-011	Plate sealer	4 strips	Room temperature

Summary

Product Description	ARG81557 Human ADAMTS1 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Human ADAMTS1 in serum, plasma (heparin) and cell culture supernatants.
Tested Reactivity	Hu
Tested Application	ELISA
Specificity	There is no detectable cross-reactivity with other relevant proteins.
Target Name	ADAMTS1
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	46.9 pg/ml
Sample Type	Serum, plasma (heparin) and cell culture supernatants.
Standard Range	93.8 - 6000 pg/ml
Sample Volume	100 µl

Precision	Intra-Assay CV: 6.3% Inter-Assay CV: 7.5%
Alternate Names	ADAMTS-1; A disintegrin and metalloproteinase with thrombospondin motifs 1; METH-1; METH1; EC 3.4.24.-; ADAM-TS1; C3-C5; ADAM-TS 1

Application Instructions


Assay Time	~ 5 hours
------------	-----------

Properties

Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	ADAMTS1
Gene Full Name	ADAM metalloproteinase with thrombospondin type 1 motif, 1
Background	This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motif) protein family. Members of the family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The protein encoded by this gene contains two disintegrin loops and three C-terminal TS motifs and has anti-angiogenic activity. The expression of this gene may be associated with various inflammatory processes as well as development of cancer cachexia. This gene is likely to be necessary for normal growth, fertility, and organ morphology and function. [provided by RefSeq, Jul 2008]
Function	Cleaves aggrecan, a cartilage proteoglycan, and may be involved in its turnover (By similarity). Has angiogenic inhibitor activity. Active metalloproteinase, which may be associated with various inflammatory processes as well as development of cancer cachexia. May play a critical role in follicular rupture. [UniProt]
Highlight	Related products: ADAMTS antibodies ; ADAMTS ELISA Kits ; New ELISA data calculation tool: Simplify the ELISA analysis by GainData
PTM	The precursor is cleaved by a furin endopeptidase. Glycosylated. Can be O-fucosylated by POFUT2 on a serine or a threonine residue found within the consensus sequence C1-X(2)-(S/T)-C2-G of the TSP type-1 repeat domains where C1 and C2 are the first and second cysteine residue of the repeat, respectively. Fucosylated repeats can then be further glycosylated by the addition of a beta-1,3-glucose residue by the glucosyltransferase, B3GALTL. Fucosylation mediates the efficient secretion of ADAMTS family members. Also can be C-glycosylated with one or two mannose molecules on tryptophan residues within the consensus sequence W-X-X-W of the TPRs, and N-glycosylated. These other glycosylations can also facilitate secretion (By similarity). [UniProt]



ARG81557 Human ADAMTS1 ELISA Kit standard curve image

ARG81557 Human ADAMTS1 ELISA Kit results of a typical standard run with optical density reading at 450 nm.