

Product datasheet

info@arigobio.com

ARG81845 Mouse MFG-E8 / Lactadherin ELISA Kit

Package: 96 wells Store at: 4°C

Component

Cat. No.	Component Name	Package	Temp
ARG81845-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG81845-002	Standard	2 X 10 ng/vial	4°C
ARG81845-003	Standard/Sample diluent	30 ml (Ready to use)	4°C
ARG81845-004	Antibody conjugate concentrate (100X)	1 vial (100 μl)	4°C
ARG81845-005	Antibody diluent buffer	12 ml (Ready to use)	4°C
ARG81845-006	HRP-Streptavidin concentrate (100X)	1 vial (100 μl)	4°C
ARG81845-007	HRP-Streptavidin diluent buffer	12 ml (Ready to use)	4°C
ARG81845-008	25X Wash buffer	20 ml	4°C
ARG81845-009	TMB substrate	10 ml (Ready to use)	4°C (Protect from light)
ARG81845-010	STOP solution	10 ml (Ready to use)	4°C
ARG81845-011	Plate sealer	4 strips	Room temperature

Summary

Product Description	ARG81845 Mouse MFG-E8 / Lactadherin ELISA Kit is an Enzyme Immunoassay kit for the quantification of Mouse MFG-E8 / Lactadherin in serum, plasma (heparin, EDTA) and cell culture supernatants.	
Tested Reactivity	Ms	
Tested Application	ELISA	
Specificity	There is no detectable cross-reactivity with other relevant proteins.	
Target Name	MFGE8 / Lactadherin	
Conjugation	on HRP	
Conjugation Note	tion Note Substrate: TMB and read at 450 nm.	
Sensitivity		
Sample Type		
Standard Range	62.5 - 4000 pg/ml	
Sample Volume	100 μΙ	

Precision Intra-Assay CV: 6.4%; Inter-Assay CV: 7.4%

Alternate Names SPAG10; MFGM; HsT19888; HMFG; Milk fat globule-EGF factor 8; MFG-E8; BA46; hP47; EDIL1;

Lactadherin; OAcGD3S; Breast epithelial antigen BA46; SED1

Application Instructions

Assay Time ~ 5 hours

Properties

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol MFGE8

Gene Full Name milk fat globule-EGF factor 8 protein

Background This gene encodes a preproprotein that is proteolytically processed to form multiple protein products.

The major encoded protein product, lactadherin, is a membrane glycoprotein that promotes

phagocytosis of apoptotic cells. This protein has also been implicated in wound healing, autoimmune disease, and cancer. Lactadherin can be further processed to form a smaller cleavage product, medin, which comprises the major protein component of aortic medial amyloid (AMA). Alternative splicing

results in multiple transcript variants. [provided by RefSeq, Jul 2015]

Function Plays an important role in the maintenance of intestinal epithelial homeostasis and the promotion of

mucosal healing. Promotes VEGF-dependent neovascularization (By similarity). Contributes to phagocytic removal of apoptotic cells in many tissues. Specific ligand for the alpha-v/beta-3 and alpha-v/beta-5 receptors. Also binds to phosphatidylserine-enriched cell surfaces in a receptor-independent manner. Zona pellucida-binding protein which may play a role in gamete interaction. Binds specifically


to rotavirus and inhibits its replication.

Medin is the main constituent of aortic medial amyloid. [UniProt]

Highlight Related products:

MFGE8 antibodies; MFGE8 ELISA Kits; New ELISA data calculation tool: Simplify the ELISA analysis by GainData

PTM Medin has a ragged N-terminus with minor species starting at Pro-264 and Gly-273. [UniProt]



ARG81845 Mouse MFG-E8 / Lactadherin ELISA Kit standard curve image $\,$

ARG81845 Mouse MFG-E8 / Lactadherin ELISA Kit results of a typical standard run with optical density reading at 450 nm.