

Product datasheet

info@arigobio.com

ARG82683 Porcine TNF alpha ELISA Kit

Package: 96 wells Store at: 4°C

Summary

Product Description ARG82683 Porcine TNF alpha ELISA Kit is an Enzyme Immunoassay kit for the quantification of Pig TNF

alpha in serum, plasma and cell culture supernatants.

Tested Reactivity Pig

Tested Application ELISA

Specificity Not react with Human TNF beta.

Not react with Human and Mouse TNF RI, TNF RII and Rat TNF alpha.

Target Name TNF alpha

Conjugation HRP

Conjugation Note Substrate: TMB and read at 450 nm.

Sensitivity 15.6 pg/ml

Sample Type Serum, plasma and cell culture supernatants.

Standard Range 31.25 - 2000 pg/ml

Sample Volume $100 \ \mu l$

Precision Intra-Assay CV: less than 9%

Inter-Assay CV: less than 9%

Alternate Names Tumor necrosis factor ligand superfamily member 2; DIF; Cachectin; ICD2; ICD1; N-terminal fragment;

TNF-a; TNFA; TNFSF2; TNF-alpha; Tumor necrosis factor; NTF

Application Instructions

Assay Time ~ 3.5 hours

Properties

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol TNF

Gene Full Name tumor necrosis factor

Function Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. It is mainly secreted by macrophages

and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under certain

www.arigobio.com arigo.nuts about antibodies 1/2

conditions it can stimulate cell proliferation and induce cell differentiation. Impairs regulatory T-cells (Treg) function in individuals with rheumatoid arthritis via FOXP3 dephosphorylation. Upregulates the expression of protein phosphatase 1 (PP1), which dephosphorylates the key 'Ser-418' residue of FOXP3, thereby inactivating FOXP3 and rendering Treg cells functionally defective (PubMed:23396208). Key mediator of cell death in the anticancer action of BCG-stimulated neutrophils in combination with DIABLO/SMAC mimetic in the RT4v6 bladder cancer cell line (PubMed:22517918, PubMed:16829952, PubMed:23396208). Induces insulin resistance in adipocytes via inhibition of insulin-induced IRS1 tyrosine phosphorylation and insulin-induced glucose uptake. Induces GKAP42 protein degradation in adipocytes which is partially responsible for TNF-induced insulin resistance (By similarity).

The TNF intracellular domain (ICD) form induces IL12 production in dendritic cells. [UniProt]

Highlight

Related products:

TNF alpha antibodies; TNF alpha ELISA Kits; TNF alpha Duos / Panels; TNF alpha recombinant proteins;

Related news:

HMGB1 in inflammation
Inflammatory Cytokines
New ELISA data calculation tool:

Simplify the ELISA analysis by GainData

PTM

The soluble form derives from the membrane form by proteolytic processing. The membrane-bound form is further proteolytically processed by SPPL2A or SPPL2B through regulated intramembrane proteolysis producing TNF intracellular domains (ICD1 and ICD2) released in the cytosol and TNF C-domain 1 and C-domain 2 secreted into the extracellular space.


The membrane form, but not the soluble form, is phosphorylated on serine residues. Dephosphorylation of the membrane form occurs by binding to soluble TNFRSF1A/TNFR1.

O-glycosylated; glycans contain galactose, N-acetylgalactosamine and N-acetylneuraminic acid. [UniProt]

Cellular Localization

Cell membrane; Single-pass type II membrane protein. Tumor necrosis factor, membrane form: Membrane; Single-pass type II membrane protein. Tumor necrosis factor, soluble form: Secreted. C-domain 1: Secreted. C-domain 2: Secreted. [UniProt]

Images



ARG82683 Porcine TNF alpha ELISA Kit standard curve image

ARG82683 Porcine TNF alpha ELISA Kit results of a typical standard run with optical density reading at 450 nm.