

## ARG83021 Mouse CCL17 / TARC ELISA Kit

Package: 96 wells  
Store at: 4°C

### Summary

Product Description	ARG83021 Mouse CCL17 / TARC ELISA Kit is an Enzyme Immunoassay kit for the quantification of Mouse CCL17 / TARC in serum, plasma (heparin), ascites, urine and cell culture supernatants.
Tested Reactivity	Ms
Tested Application	ELISA
Specificity	Not cross-reacts with: Human IL2, IL3, IL4, IL6, IL10, CCL17 / TRAC, MCP1, MCP2. Mouse IL2, IL3, IL4, IL6, IL10, TNFRI, TNFR2, VEGF, RANTES.
Target Name	CCL17
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	15.6 pg/ml
Sample Type	Serum, plasma (heparin), ascites, urine and cell culture supernatants.
Standard Range	31.25 - 2000 pg/ml
Sample Volume	100 µl
Precision	Intra-Assay CV: less than 10% Inter-Assay CV: less than 10%
Alternate Names	CCL17, C C Motif Chemokine Ligand 17, TARC, ABCD 2, SCYA17, Small Inducible Cytokine Subfamily A (Cys-Cys), Member 17, Thymus And Activation-Regulated Chemokine, Chemokine (C C Motif) Ligand 17, Small Inducible Cytokine A17, C C Motif Chemokine 17, CC Chemokine TARC, T Cell Directed CC Chemokine, A-152E5.3

### Application Instructions

Assay Time	~ 4 hours
------------	-----------

### Properties


Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

Gene Symbol	CCL17
Gene Full Name	C-C Motif Chemokine Ligand 17

Background	This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for T lymphocytes, but not monocytes or granulocytes. The product of this gene binds to chemokine receptors CCR4 and CCR8. This chemokine plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells. [provided by RefSeq, Sep 2014]
Function	Chemotactic factor for T-lymphocytes but not monocytes or granulocytes. May play a role in T-cell development in thymus and in trafficking and activation of mature T-cells. Binds to CCR4. [UniProt]
Highlight	

Images



ARG83021 Mouse CCL17 / TARC ELISA Kit standard curve image

ARG83021 Mouse CCL17 / TARC ELISA Kit results of standard run with optical density reading at 450 nm