

Product datasheet

info@arigobio.com

ARG83033 Human FGF19 ELISA Kit

Package: 96 wells Store at: 4°C

Summary

Product Description ARG83033 Human FGF19 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Human

FGF19 in serum, plasma (heparin), ascites, urine and cell culture supernatants.

Tested Reactivity Hu

Tested Application ELISA

Specificity Not cross-reacts with:

Human bFGF, FGF-21, FGF R1 alpha, FGF-R1 beta, FGF-R2 alpha, FGF-R2 beta.

Mouse FGF15, FGF21, FGF23, FGFR1 alpha, FGFR1 beta. Rat FGF15, FGF21, FGF23, FGFR1 alpha, FGFR1 beta.

Target Name FGF19

Conjugation HRP

Conjugation Note Substrate: TMB and read at 450 nm.

Sensitivity 15.6 pg/ml

Sample Type Serum, plasma (heparin), ascites, urine and cell culture supernatants.

Standard Range 31.3 - 2000 pg/ml

Sample Volume $$100~\mu l$$

Precision Intra-Assay CV: less than 10%

Inter-Assay CV: less than 10%

Alternate Names FGF19, FGF 19, Fibroblast Growth Factor 19

Application Instructions

Assay Time ~ 4 hours

Properties

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol FGF19

Gene Full Name Fibroblast Growth Factor 19

Background The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family

members possess broad mitogenic and cell survival activities, and are involved in a variety of biological

www.arigobio.com arigo.nuts about antibodies 1/2

processes including embryonic development cell growth, morphogenesis, tissue repair, tumor growth and invasion. This growth factor is a high affinity, heparin dependent ligand for FGFR4. Expression of this gene was detected only in fetal but not adult brain tissue. Synergistic interaction of the chick homolog and Wnt-8c has been shown to be required for initiation of inner ear development. [provided by RefSeq, Jul 2008]

Function Involved in the suppression of bile acid biosynthesis through down-regulation of CYP7A1 expression,

 $following\ positive\ regulation\ of\ the\ JNK\ and\ ERK1/2\ cascades.\ Stimulates\ glucose\ uptake\ in\ adipocytes.$

Activity requires the presence of KLB and FGFR4. [UniProt]

Highlight Related products:


FGF19 antibodies; FGF19 ELISA Kits; FGF19 recombinant proteins;

New ELISA data calculation tool: Simplify the ELISA analysis by GainData

PTM Disulfide bond. [UniProt]

Cellular Localization Secreted. [UniProt]

Images



ARG83033 Human FGF19 ELISA Kit standard curve image

ARG83033 Human FGF19 ELISA Kit results of standard run with optical density reading at 450 nm