

ARG83168 Rabbit N Cadherin ELISA Kit

Package: 96 wells
Store at: 4°C

Summary

Product Description	ARG83168 Rabbit N Cadherin ELISA Kit is an Enzyme Immunoassay kit for the quantification of Rabbit N Cadherin in Serum, Plasma and Cell culture supernatants.
Tested Reactivity	Rb
Tested Application	ELISA
Specificity	There is no detectable cross-reactivity with other relevant proteins.
Target Name	N Cadherin
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	25 pg/ml
Detection Range	156 pg/ml - 10,000 pg/ml
Sample Type	Serum, Plasma and Cell culture supernatants
Precision	Intra-Assay CV: 5.2% Inter-Assay CV: 5.8%
Alternate Names	Neural cadherin; N-cadherin; CDw325; CDHN; CD antigen CD325; NCAD; Cadherin-2; CD325

Application Instructions

Assay Time	~ 5 hours
------------	-----------

Properties

Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	CDH2
Gene Full Name	cadherin 2, type 1, N-cadherin (neuronal)
Background	N Cadherin is a classical cadherin and member of the cadherin superfamily. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein is proteolytically processed to generate a calcium-dependent cell adhesion molecule and glycoprotein. This protein plays a role in the establishment of left-right asymmetry, development of the nervous system and the formation of cartilage and bone. [provided by RefSeq, Nov 2015]
Function	N Cadherin is a calcium-dependent cell adhesion protein; preferentially mediates homotypic cell-cell

adhesion by dimerization with a CDH2 chain from another cell. Cadherins may thus contribute to the sorting of heterogeneous cell types. Acts as a regulator of neural stem cells quiescence by mediating anchorage of neural stem cells to ependymocytes in the adult subependymal zone: upon cleavage by MMP24, CDH2-mediated anchorage is affected, leading to modulate neural stem cell quiescence. CDH2 may be involved in neuronal recognition mechanism. In hippocampal neurons, may regulate dendritic spine density. [UniProt]

Highlight

Related products:

[Cadherin antibodies](#); [Cadherin ELISA Kits](#); [Cadherin Duos / Panels](#);

New ELISA data calculation tool:


[Simplify the ELISA analysis by GainData](#)

PTM

Cleaved by MMP24. Ectodomain cleavage leads to the generation of a soluble 90 kDa amino-terminal soluble fragment and a 45 kDa membrane-bound carboxy-terminal fragment 1 (CTF1), which is further cleaved by gamma-secretase into a 35 kDa. Cleavage in neural stem cells by MMP24 affects CDH2-mediated anchorage of neural stem cells to ependymocytes in the adult subependymal zone, leading to modulate neural stem cell quiescence (By similarity).

May be phosphorylated by OBSCN. [UniProt]

Images



ARG83168 Rabbit N Cadherin ELISA Kit standard curve image

ARG83168 Rabbit N Cadherin ELISA Kit results of a typical standard run with optical density reading at 450 nm.